

AARC Clinical Practice Guideline

Long-Term Invasive Mechanical Ventilation in the Home —2007 Revision & Update

HIMV 1.0 PROCEDURE

The application of invasive mechanical ventilation and care of the patient-ventilator system in the home, as ordered by a physician.

HIMV 2.0 DESCRIPTION/DEFINITION

Mechanical ventilation may be defined as a life support system designed to replace or support normal ventilatory lung function. Ventilator dependence is caused by an imbalance between ventilatory capacity and demand. A ventilator-assisted individual (VAI) may require mechanical aid for breathing to augment or replace spontaneous ventilatory efforts to achieve medical stability or to maintain life. The patient eligible for invasive long-term mechanical ventilation in the home (HIMV) requires a tracheostomy tube for ventilatory support, but no longer requires intensive medical and monitoring services.¹⁻⁶ This guideline refers to patients ventilated by positive pressure via a tracheostomy tube in the home.

2.1 The goals of HIMV are

- 2.1.1 To sustain and extend life¹⁻⁶
- 2.1.2 To enhance the quality of life¹⁻¹¹
- 2.1.3 To reduce morbidity^{1-6,12-15}
- 2.1.4 To improve or sustain physical and psychological function of all VAIs and to enhance growth and development in pediatric VAIs¹⁻¹¹
- 2.1.5 To provide cost-effective care

HIMV 3.0 SETTING

The setting is the home, which for the purposes of this guideline may be the patient's home, a foster home, or a group-living environment^{1-6,16-19}

HIMV 4.0 INDICATIONS

4.1 Patients requiring invasive long-term ventilatory support have demonstrated

- 4.1.1 An inability to be completely weaned from invasive ventilatory support or

4.1.2 A progression of disease etiology that requires increasing ventilatory support.

4.2 Conditions that met these criteria may include but are not limited to ventilatory muscle disorders, alveolar hypoventilation syndrome, primary respiratory disorders, obstructive lung diseases, restrictive lung diseases, and cardiac disorders, including congenital anomalies^{1-6,16-20}

HIMV 5.0 CONTRAINDICATIONS

Contraindications to HIMV include:

5.1 The presence of a physiologically unstable medical condition requiring higher level of care or resources than available in the home¹⁻⁶ Examples of indicators of a medical condition too unstable for the home and long-term care setting are:

5.1.1 F_{IO_2} requirement $> 0.40^{1-6}$

5.1.2 $PEEP > 10 \text{ cm H}_2\text{O}^{1-6}$

5.1.3 Need for continuous invasive monitoring in adult patients¹⁻⁶

5.1.4 Lack of mature tracheostomy

5.2 Patient's choice not to receive home mechanical ventilation^{1-6,20-24}

5.3 Lack of an appropriate discharge plan¹⁻⁶

5.4 Unsafe physical environment as determined by the patient's discharge planning team¹⁻⁶

5.4.1 Presence of fire, health or safety hazards including unsanitary conditions¹⁻⁶

5.4.2 Inadequate basic utilities (such as heat, air conditioning, electricity including adequate amperage and grounded outlets)¹⁻⁶

5.5 Inadequate resources for care in the home

5.5.1 Financial^{1-6,25-28}

5.5.2 Personnel

5.5.2.1 Inadequate medical follow-up¹⁻⁶

5.5.2.2 Inability of VAI to care for self, if no caregiver is available¹⁻⁶

5.5.2.3 Inadequate respite care for caregivers^{21-23,29,30}

5.5.2.4 Inadequate numbers of competent caregivers¹⁻⁶ A minimum of two competent caregivers are required.

HIMV 6.0 HAZARDS AND COMPLICATIONS

6.1 Deterioration or acute change in clinical status of VAI. Although ventilator-associated complications in the home are poorly documented, experience in other sites can be extrapolated. The following may cause death or require rehospitalization for acute treatment.

6.1.1 Medical: Hypocapnia, respiratory alkalosis hypercapnia, respiratory acidosis, hypoxemia, barotraumas, seizures, hemodynamic instability, airway complications (stomal or tracheal infection, mucus plugging, tracheal erosion, or stenosis), respiratory infection (tracheobronchitis, pneumonia, bronchospasm, exacerbation of underlying disease, or natural course of the disease^{1-6,14})

6.1.2 Equipment-related: Failure of the ventilator, malfunction of equipment, inadequate warming, and humidification of the inspired gases, inadvertent changes in ventilator settings, accidental disconnection from ventilator, accidental decannulation^{1-6,31-36}

6.1.3 Psychosocial: Depression, anxiety, loss of resources (caregiver or financial), detrimental change in family structure or coping capacity^{1-6,21-24,29,30,37,38}

HIMV 7.0 LIMITATIONS

In the home care setting, making and implementing changes in the plan of care may take longer than in a health care facility.

HIMV 8.0 ASSESSMENT OF NEED

8.1 Determination that indications are present and contraindications are absent

8.2 Determination that the goals listed in 2.1 can be met in the home

8.3 Determination that no continued need exists for higher level of services

8.4 Determination that frequent changes in the plan of care will not be needed

HIMV 9.0 ASSESSMENT OF OUTCOME

At least the following aspects of patient management and condition should be evaluated periodically as long as the patient receives HIMV

9.1 Implementation and adherence to the plan of care

9.2 Quality of life

9.3 Patient satisfaction

9.4 Resource utilization

9.5 Growth and development in the pediatric patient

9.6 Change in prognosis

9.7 Unanticipated morbidity, including need for higher level site of care

9.8 Unanticipated mortality

HIMV 10.0 RESOURCES

10.1 Equipment

10.1.1 Ventilator(s)—Choice should be based on patient's clinical need. Patient's medical needs may dictate that more than one ventilator be provided.¹⁻⁶

10.1.1.1 Ventilators chosen for home care must be dependable and easy for the intended caregivers to operate; small size and lightweight are desirable.

10.1.1.2 Mobility is frequently an essential element of the plan of care of the patient. The mechanical ventilator system chosen for such a patient should allow mobility.

10.1.2 With portable, volume-cycled ventilators, use of the SIMV mode increases work of breathing¹⁻⁶

10.1.3 Complex and non-portable components are not recommended for HIMV but may be used to meet the needs of certain patients¹⁻⁶

10.1.3.1 Ventilators powered by external compressed gas sources are less desirable¹⁻⁶

10.1.3.2 A second ventilator should be provided for

10.1.3.2.1 Patients who cannot maintain spontaneous ventilation for 4 or more consecutive hours¹⁻⁶

10.1.3.2.2 Patients who live in an area where a replacement ventilator cannot be provided within 2 hours¹⁻⁶

- 10.1.3.2.3** Patients who require mechanical ventilation during mobility as prescribed in their plan of care
- 10.1.4** Preventive maintenance should be provided at the frequency recommended under manufacturer guidelines
- 10.1.5** An adequate power source must be available to operate the ventilator consistent with patient needs.¹⁻⁶ This may be supplied by one or more of the following methods
- 10.1.5.1** Alternating current (AC) is the primary power source for most long-term care ventilators. Emergency AC power should be available in the long-term care facility.
 - 10.1.5.2** Direct current (DC) by external battery may be used to allow mobility and as an emergency power source. The internal battery of the ventilator should be used only for short-term use. It should not be used as a primary source of power.
 - 10.1.5.3** A portable generator may be recommended for the VAI if frequent power outages occur or if the home is in a remote location
- 10.1.6 Alarms**
- 10.1.6.1** A patient-disconnect (eg, low-pressure or low-exhaled-volume) and a high-pressure alarm are essential.
 - 10.1.6.2** If patient disconnection is likely to produce a serious adverse effect, a remote alarm and a secondary alarm may be indicated. A secondary alarm may be based on chest-wall impedance and cardiac activity, exhaled volume, end-tidal CO₂, or pulse oximetry with alarm capabilities^{3,5}
 - 10.1.6.3** Audible alarms must be loud enough to be heard by caregivers in all areas of the home.
- 10.1.7** Humidification systems are essential for invasive mechanical ventilation. The type of system used is determined by the patient's medical needs and the patient's need for mobility. It may be appropriate for the patient to use more than one type of system, based on those needs.¹⁻⁵
- 10.1.7.1** Heated humidifier (temperature probes should be provided)
- 10.1.7.2** Heat - moisture exchanger (HME) can be used during transport and to enhance mobility and may be used in lieu of a heated humidifier if the HME is determined to meet the patient's medical needs^{1-3,5,6}
- 10.1.8** Ventilator circuit and accessories as medically indicated
- 10.1.9** Self-inflating resuscitation bag with tracheostomy attachments, oxygen port if oxygen is prescribed, and mask of appropriate size^{1-3,5,6}
- 10.1.10** Replacement tracheostomy tube of appropriate size, plus a tube one size smaller should be available at all times
- 10.1.11** Suction equipment^{1-3,5,6} including a battery-powered aspirator for patients who leave the home or when indicated as an alternate source in the event of a power failure
- 10.1.12** Supplemental oxygen as medically indicated^{1-3,5,6}
- 10.1.13** VAI must have an adequate means of communicating their needs/desires and have the means to summon help from their caregivers in the case of emergency^{3,5}
- 10.1.14** VAI and caregivers must have functioning phone lines so that they can contact and be contacted by medical personnel in the case of emergency
- 10.2 Personnel**
- 10.2.1** Health care professionals capable of providing direct patient care and possessing demonstrated competencies to monitor and assess both the patient and equipment are essential. Health care professionals should be credentialed (RRT, CRT, RN) and/or licensed practitioners with documented knowledge and demonstrated competencies so as to:
 - 10.2.1.1** Understand the patient's disease, plan of care, goals, and the limitations of invasive mechanical ventilation^{2,3,5,6}
 - 10.2.1.2** Assess patient's response to invasive mechanical ventilation

10.2.1.3 Make recommendations for changes in respiratory management of patient, including weaning as necessary^{2,3,5,6}

10.2.1.4 Train and monitor lay caregivers

10.2.1.5 Monitor patient's ongoing ventilatory status

10.2.1.6 Communicate results of assessment to the health care team^{2,3,5,6}

10.2.2 Lay caregivers (family members, personal care attendants, non-credentialed health care personnel such as nurse's aides) can be taught tasks and techniques of care for a specific VAI. Appropriately trained lay caregivers must demonstrate competency in:

10.2.2.1 Proper set up, use, troubleshooting, and routine maintenance of the equipment and supplies^{2,3,5,6,21,23,24,29,30,37-41}

10.2.2.2 Identification of adverse patient response to invasive mechanical ventilation^{2,3,5,6}

10.2.2.3 Response to the hazards of invasive mechanical ventilation^{2,3,5,6}

10.2.2.4 Response to emergencies^{2,3,5,6} such as

10.2.2.4.1 Power failure

10.2.2.4.2 Acute life threatening events such as accidental decannulation or medical deterioration of the patient

10.2.2.4.3 Failure of the equipment or supplies

10.2.2.5 Appropriate infection control procedures^{2,3,5,6}

10.2.2.6 Use and application of any additional techniques required for ongoing care of the VAI, such as suctioning and the use of ancillary equipment^{2,3,5,6}

10.3 Finances: HIMV can only be instituted and maintained with adequate financial resources to provide the necessary equipment and personnel to manage the patient's care^{2,3,5,6,25-28}

HIMV 11.0 MONITORING

The frequency of monitoring should be determined by the ongoing individualized care plan and be based upon the patient's current medical condition.

The ventilator settings, proper function of equipment, and the patient's physical condition should be monitored and verified: with each initiation of invasive ventilation to the patient, including altering the source of ventilation, as from one ventilator or resuscitation bag to another ventilator; with each ventilator setting change; after moving the patient (eg, from the bed to a chair); on a regular basis as specified by individualized plan of care.³ All caregivers, both professional and appropriately trained lay caregivers, should follow the care plan and implement the monitoring that has been prescribed. After being trained and evaluated on their level of knowledge and ability to respond to the VAI clinical response to each intervention, lay caregivers with documented competency may operate, perform routine maintenance tasks, monitor equipment, and perform personal care required by the VAI.

11.1 After completing training, demonstrating competency and if directed in the VAI's plan of care, the lay caregivers should monitor the following

11.1.1 Patient's physical condition (may include the following: respiratory rate, heart rate, color changes, chest excursion, diaphoresis and lethargy, blood pressure, body temperature)

11.1.2 Ventilator settings. The frequency at which alarms and settings are to be checked should be specified in the plan of care.

11.1.2.1 Peak pressures

11.1.2.2 Preset tidal volume or preset pressure control

11.1.2.3 Frequency of ventilator breaths

11.1.2.4 Verification of oxygen concentration setting or flow rate of oxygen bled into the ventilator system

11.1.2.5 PEEP level (if applicable)

11.1.2.6 Appropriate humidification of inspired gases

11.1.2.7 Temperature of inspired gases (if applicable)

11.1.2.8 Heat-moisture exchanger (HME) function (if applicable)

11.1.3 Equipment function^{3,5}

11.1.3.1 Appropriate configuration of ventilator circuit^{3,5}

11.1.3.2 Alarm function

11.1.3.3 Cleanliness of filter(s)—according to manufacturer's recommendation

11.1.3.4 Battery power level(s)—both internal and external

11.1.3.5 Overall condition of all equipment

11.1.3.6 Self-inflating manual resuscitator—cleanliness and function

11.2 Health care professionals should perform a thorough, comprehensive assessment of the patient and the patient-ventilator system on a regular basis as prescribed by the plan of care. In addition to the variables listed in 11.1.1–11.1.3.6, the health care professional should implement, monitor, and assess results of other interventions as indicated by the clinical situation and anticipated in the care plan.

11.2.1 Pulse oximetry—should be used to assess patients requiring a change in prescribed oxygen levels or in patients with a suspected change in condition^{3,5}

11.2.1.1 A physician's order for pulse oximetry must be obtained before oximetry testing is performed

11.2.2 End-tidal CO₂—may be useful for establishing trends in CO₂ levels^{3,20}

11.2.2.1 A physician's order for end tidal CO₂ monitoring must be obtained before end tidal CO₂ monitoring is performed

11.2.3 Ventilator settings

11.2.4 Exhaled tidal volume

11.2.5 Analysis of fraction of inspired oxygen

11.3 Health care professionals are also responsible for maintaining interdisciplinary communication concerning the plan of care

11.4 Health care professionals should integrate respiratory plan of care into the patient's total care plan.^{2,3,5,6} Plan of care should include

11.4.1 All aspects of patient's respiratory care^{2,3,5,6}

11.4.2 Ongoing assessment and education of the caregivers involved

HIMV 12.0 FREQUENCY:

12.1 The frequency of ventilation (and the patient's ventilator-free time) is dictated by the patient's physiologic needs and is deter-

mined in consultation with the patient's physician.

12.2 The frequency of assessment of the VAI and the patient-ventilator system must be noted in the evolving total care plan as determined by the health care team, in conjunction with the VAI and their caregivers.

HIMV 13.0 INFECTION CONTROL

13.1 Both professional and lay caregivers should be aware of the potential for transmission of both chronic and acute infection from patient to caregiver and from caregiver to patient and should take the steps necessary to avoid that transmission. Aspects of avoidance include

13.1.1 Careful hand cleansing and barrier protection when appropriate

13.1.2 Careful disposal of medical waste

13.1.3 Maximizing protection of patient, family, and caregivers (eg, influenza immunization) and minimizing exposure to persons with acute infections (eg, limiting visitors with upper respiratory infections)

13.2 Evidence is lacking to support an optimal plan for changing and processing ventilator circuits and ancillary equipment in the home. The standard of care in the home is that ventilator circuits need not be changed more often than once each week. However, CDC guidelines and studies from institutional settings^{42–44} suggest that ventilator circuits need only be changed when visibly soiled.

Revised by

Respiratory Home Care Focus Group:

Joan Kohorst MA RRT

Patricia Blakely RRT

Claude Dockter RRT

William Pruitt MBA RRT CPFT AE-C

Originally published in Respiratory Care
1995;40(12):1313-1320.

REFERENCES

- Pilbeam S, Gramlich T. Long-term ventilation. In: Pilbeam S, Cairo J. Mechanical ventilation: physiological and clinical applications. St Louis: Mosby Elsevier; 2006: 473-507.

2. Ambrosino N, Vianello A. Where to perform long-term ventilation. *Respir Care Clin N Am* 2002;8(3):463-478.
3. Make BJ, Hill NS, Goldberg AI, Bach JR, Criner GJ, Dunne PE, et al. Mechanical ventilation beyond the intensive care unit: report of a consensus conference of the American College of Chest Physicians. *Chest* 1998;113(5 Suppl):289S-344S.
4. Stuart M, Weinrich M. Protecting the most vulnerable: home mechanical ventilation as a case study in disability and medical care: report from an NIH conference. *Neurorehabil Neural Repair* 2001;15(3):159-166.
5. Kohorst J. Transitioning the ventilator-dependent patient from the hospital to home. *Medscape Pulmonary Medicine* 2005. [Internet]. Available at: <http://www.medscape.com/viewarticle/514735>. (Last accessed Feb 20, 2007.)
6. Gilmartin M. Transition from the intensive care unit to home: patient selection and discharge planning. *Respir Care* 1994;39(5):456-480.
7. Markstrom A, Sundell K, Lysdahl M, Andersson G, Schedin U, Klang B. Quality-of-life evaluation of patients with neuromuscular and skeletal diseases treated with noninvasive and invasive home mechanical ventilation. *Chest* 2002;122(5):1695-1700.
8. Goldstein RS, Psek JA, Gort EH. Home mechanical ventilation: demographics and user perspectives. *Chest* 1995;108(6):1581-1586.
9. Brooks D, King A, Tonack M, Simson H, Gould M, Goldstein R. User perspectives on issues that influence the quality of daily life of ventilator-assisted individuals with neuromuscular disorders. *Can Respir J* 2004;11(8):547-554.
10. Douglas SL, Daly BJ, Gordon N, Brennan PF. Survival and quality of life: short-term versus long-term ventilator patients. *Crit Care Med* 2002;30(12):2655-2662.
11. Chelluri L, Im KA, Belle SH, Schulz R, Rotondi AJ, Donahoe MP, et al. Long-term mortality and quality of life after prolonged mechanical ventilation. *Crit Care Med* 2004;32(1):61-69.
12. Douglas SL, Daly BJ, Brennan PF, Harris S, Nochomovitz M, Dyer MA. Outcomes of long-term ventilator patients: a descriptive study. *Am J Crit Care* 1997;6(2):99-105.
13. Im K, Belle SH, Schulz R, Mendelsohn AB, Chelluri L; QOL-MV Investigators. Prevalence and outcomes of caregiving after prolonged (\geq 48 hours) mechanical ventilation in the ICU. *Chest* 2004;125(2):597-606.
14. Douglas SL, Daly BJ, Brennan PF, Gordon NH, Uthis P. Hospital readmission among long-term ventilator patients. *Chest* 2001;120(4):1278-1286.
15. Douglas SL, Daly BJ. Caregivers of long-term ventilator patients: physical and psychological outcomes. *Chest* 2003;123(4):1073-1081.
16. Make BJ, Gilmartin ME. Mechanical ventilation in the home. *Crit Care Clin* 1990;6(3):785-796.
17. Rouault S. [Education of the families of patients receiving domiciliary ventilation.] *Rev Mal Respir* 2005;22(3):461-465. (Article in French)
18. Gonzalez J, Macey J, Cracco C, Similowski T, Derenne JP. [Initiation and management of home mechanical ventilation.] *Rev Mal Respir* 2004;21(4 pt 1):783-790. (Article in French)
19. Glass C, Grap MJ, Battle G. Preparing the patient and family for home mechanical ventilation. *Medsurg Nurs* 1999;8(2):99-101,104-107.
20. MacIntyre N, Cook D, Ely E Jr, Epstein S, Fink J, Heffner JE, et al. Evidence-based guidelines for weaning and discontinuing ventilatory support: a collective task force facilitated by the American College of Chest Physicians; the American Association for Respiratory Care; and the American College of Critical Care Medicine. *Chest* 2001;120(6 Suppl):375S-395S.
21. Lehoux P, Saint-Arnaud J, Richard L. The use of technology at home: what patient manuals say and sell vs. what patients face and fear. *Sociol Health Illn* 2004;26(5):617-644.
22. Lindahl B, Sandman PO, Rasmussen BH. On becoming dependent on home mechanical ventilation. *J Adv Nurs* 2005;49(1):33-42.
23. Lindahl B, Sandman PO, Rasmussen BH, On being dependent on home mechanical ventilation: depictions of patients' experiences over time. *Qual Health Res* 2006;16(7):881-901.
24. Beach SR, Schulz R, Williamson GM, Miller LS, Weiner MF, Lance CE. Risk factors for potentially harmful informal caregiver behavior. *J Am Geriatr Soc* 2005;53(2):255-261.
25. Ely EW, Baker AM, Evans GW, Haponik EF. The distribution of costs of care in mechanically ventilated patients with chronic obstructive pulmonary disease. *Crit Care Med* 2000;28(2):408-413.
26. Guber A, Morris E, Chen B, Israel S. First experience with the home-care management system for the respiratory patients in Israel. *Isr Med Assoc J* 2002;4(6):418-420.
27. Lewarski JS. Home mechanical ventilation update. *AARC Times* 2005;29(9):4-8.
28. Kurek CJ, Cohen IL, Lambrinos J, Minatoya K, Booth FV, Chalfin DB. Clinical and economic outcome of patients undergoing tracheostomy for prolonged mechanical ventilation in New York state during 1993: analysis of 6,353 cases under diagnosis-related group 483. *Crit Care Med* 1997;25(6):983-988.
29. Parra MM. Nursing and respite care services for ventilator-assisted children. *Caring* 2003;22(5):6-9.
30. Carnevale FA, Alexander E, Davis M, Rennick J, Troini R. Daily living with distress and enrichment: the moral experience of families with ventilator-assisted children at home. *Pediatrics* 2006;117(1):e48-e60.
31. Amin RS, Fitton CM. Tracheostomy and home ventilation in children. *Semin Neonatol* 2003;8(2):127-135.
32. Boyer A, Thiery G, Lasry S, Pigne E, Salah A, de Lassence A, et al. Long-term mechanical ventilation with hygroscopic heat and moisture exchangers used for 48 hours: a prospective clinical, hygrometric, and bacteriologic study. *Crit Care Med* 2003;31(3):823-829.

33. Farre R, Navajas D, Prats E, Martí S, Guell R, Montserrat JM, et al. Performance of mechanical ventilators at the patient's home: a multicentre quality control study. *Thorax* 2006;61(5):400-404.
34. Farre R, Giro E, Casolive V, Navajas D, Escarrabill J. Quality control of mechanical ventilation at the patient's home. *Intensive Care Med* 2003;29(3):484-486.
35. Srinivasan S, Doty SM, White TR, Segura VH, Jansen MT, Davidson Ward SL, Keens TG. Frequency, causes and outcome of home ventilator failure. *Chest* 1998;114(5):1363-1367.
36. Branson RD, Johannigman JA. What is the evidence base for the newer ventilation modes? *Respir Care* 2004;49(7):742-760.
37. Wang KW, Barnard A. Technology-dependent children and their families: a review. *J Adv Nurs* 2004;45(1):36-46.
38. Heaton J, Noyes J, Sloper P, Shah R. Families' experience of caring for technology-dependent children: a temporal perspective. *Health Soc Care Community* 2005;13(5):441-450.
39. Corlett J, Twycross A. Negotiation of parental roles within family-centered care: a review of the research. *J Clin Nurs* 2006;15(10):1308-1316.
40. Kirk S, Glendinning C, Callery P. Parent or nurse? The experience of being the parent of a technology-dependent child. *J Adv Nurs* 2005;51(5):456-464.
41. Kirk S. Negotiating lay and professional roles in the care of children with complex health care needs. *J Adv Nurs* 2001;34(5):593-602.
42. Hess D, Burns E, Romagnoli D, Kacmarek RM. Weekly ventilator circuit changes: a strategy to reduce costs without affecting pneumonia rates. *Anesthesiology* 1995;82(4):903-911.
43. Kollef MH, Shapiro SD, Fraser VJ, Silver P, Murphy DM, Trovillion E, et al. Mechanical ventilation with or without 7-day circuit changes: a randomized controlled trial. *Ann Intern Med* 1995;123(3):168-174.
44. American Association for Respiratory Care. AARC Evidence-Based Clinical Practice Guideline: Care of the ventilator circuit and its relation to ventilator-associated pneumonia. *Respir Care* 2003;48(9):869-879.

Interested persons may photocopy these Guidelines for noncommercial purposes of scientific or educational advancement. Please credit AARC and RESPIRATORY CARE Journal.

All of the AARC CPGs may be downloaded at no charge from

<http://www.RCJournal.com/>